



CURESUPPORT HOLDING B.V.
ZUTPHENSEWEG 55, 7418 AH
DEVENTER, THE NETHERLANDS.

BIOAVAILABILITY STUDY OF
CURESUPPORT LIPOSOMAL
CURCUMINE.

STUDY NO: CT-0007-17

THIS BIOAVAILABILITY STUDY
IS PERFORMED BY BIO AGILE
THERAPEUT.



BIOAVAILABILITY
STUDY OF

LIPOSOMAL VITAMIN C

THIS BIOAVAILABILITY STUDY
IS PERFORMED BY BIO
AGILE THERAPEUT

CURESUPPORT
IT'S ALL ABOUT ABSORPTION

BIOAVAILABILITY STUDY OF LIPOSOMAL VITAMIN C

OBJECTIVE OF THE STUDY

To evaluate the oral bioavailability of Liposomal Vitamin C 5ml/1000mg of Curesupport Holding B.V. Zutphenseweg 55, Deventer and Non Liposomal Vitamin C 5ml/1000mg (Reference) Of Curesupport Holding B.V. Zutphenseweg 55, Deventer. In healthy, adult, human subjects under fasting condition.

NUMBER OF SUBJECTS

Number of subjects planned: 24
Number of subjects completed the study and analyzed: 24

TESTED PRODUCT

Product name:
Liposomal Vitamin C 5ml/ 1000mg

Manufactured by:
Curesupport Holding B.V.
Zutphenseweg 55, 7418 AH,
Deventer, The Netherlands.

Batch No: 171106

REFERENCE PRODUCT

Product name:
Non Liposomal Curcumin 500 mg (250 mg curcumin) per dosage of 25 ml

Manufactured by:
Curesupport Holding B.V.
Zutphenseweg 55, 7418 AH,
Deventer, The Netherlands.

Lot No: 171164


DOSE AND MODE OF ADMINISTRATION

A single oral dose, 5 ml of test product (T) or Reference product was administered by using syringe (as per randomization schedule) orally, after administration of Investigational Medicinal products, syringe was rinsed with 240 ml of drinking water, at ambient temperature, to the subjects in sitting posture under fasting condition. Treatment administration was separated by wash out period 7 days.

CONCLUSION

Based on the results obtained in this study, the test product (T).

Liposomal Vitamin C 5ml/1000mg of Curesupport Holding B.V. Zutphenseweg 55, Deventer are found to be at least 6.6 times more bioavailable than published reports (on healthy, adult, human subjects under fasting conditions) of the competition.



MEANS VS TIME

■ R
■ T

N=24

RESULTS

GEOMETRIC MEAN OF REFERENCE PRODUCT (R) AND TEST PRODUCT (T) FOR ASCORBIC ACID (VITAMIN C)

PHARMACOKINETIC PARAMETER	N	REFERENCE PRODUCT (R)	TEST PRODUCT (T)
C _{max} (mg/dL)	24	2.1695	5.2386
AUC _{0-t} (mg*hr/dL)	24	31.5286	55.8627
AUC _{0-∞} (mg*hr/dL)	24	57.1165	78.9010
t _{max} (hr)	24	3.42	3.51
K _{el} (1/hr)	24	0.0365	0.0559
t _{1/2} (hr)	24	18.9995	12.3895

STATISTICAL RESULTS OF TEST PRODUCT (T) VERSUS REFERENCE PRODUCT (R) FOR VITAMIN C

PHARMACOKINETIC PARAMETER	GEOMETRIC LEAST SQUARE MEAN		INTRA SUBJECT CV (%) (T VS. R)	T/R RATIO(%)
	TEST PRODUCT (T)	REFERENCE PRODUCT (R)		
C _{max} (mg/dL)	5.2386	2.1695	8.19%	241.47%
AUC _{0-t} (mg*hr/dL)	55.8627	31.5286	13.32%	177.18%
AUC _{0-∞} (mg*hr/dL)	78.9010	57.1165	23.66%	138.14%

ORAL BIOAVAILABILITY VALUE FOR LIPOSOMAL VITAMIN C

PHARMACOKINETIC PARAMETER	TEST PRODUCT (MEAN ± STANDARD DEVIATION)	REFERENCE PRODUCT (MEAN ± STANDARD DEVIATION)	ORAL BIOAVAILABILITY VALUE (AUC _{0-t} OF TEST / AUC _{0-t} OF REFERENCE)
AUC _{0-t} (mg.hr/dL)	(56.3218 ± 7.3274)	(31.9704 ± 7.3274)	1.76